

10/25/00
U.S. PTO

10/25/00

PTO/SB/05 (08-00)

Approved for use through 10/31/2002. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Please type a plus sign (+) inside this box → +

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

09/696500 PRO
10/25/00

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

1. Fee Transmittal Form (e.g., PTO/SB/17)
(Submit an original and a duplicate for fee processing)
2. Applicant claims small entity status.
See 37 CFR 1.27.
3. Specification [Total Pages 28]
(preferred arrangement set forth below)
 - Descriptive title of the invention
 - Cross Reference to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to sequence listing, a table, or a computer program listing appendix
 - Background of the Invention
 - Brief Summary of the Invention
 - Brief Description of the Drawings *(if filed)*
 - Detailed Description
 - Claim(s)
 - Abstract of the Disclosure
4. Drawing(s) (35 U.S.C. 113) [Total Sheets 2]
5. Oath or Declaration [Total Pages 2]
 a. Newly executed (original or copy)
 b. Copy from a prior application (37 CFR 1.63 (d))
(for continuation/divisional with Box 17 completed)
 - i. **DELETION OF INVENTOR(S)**
 Signed statement attached deleting inventor(s)
 named in the prior application, see 37 CFR
 1.63(d)(2) and 1.33(b).
6. Application Data Sheet. See 37 CFR 1.76

17. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment, or in an Application Data Sheet under 37 CFR 1.76:

Continuation Divisional Continuation-in-part (CIP)

of prior application No. _____ / _____

Prior application information: Examiner _____

Group / Art Unit. _____

For CONTINUATION OR DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 5b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

18. CORRESPONDENCE ADDRESS

Customer Number or Bar Code Label



or Correspondence address below

Name	Heidi S. Nebel				
Address	Zarley, McKee, Thomte, Voorhees & Sease				
	801 Grand Avenue - Suite 3200				
City	Des Moines	State	IA	Zip Code	50309-2721
Country	U.S.A.	Telephone	515-288-3667	Fax	515-288-1338

Name (Print/Type)	Heidi S. Nebel	Registration No. (Attorney/Agent)	37,719
Signature			
	Date	10/25/2000	

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

FEE TRANSMITTAL for FY 2001

Patent fees are subject to annual revision.

TOTAL AMOUNT OF PAYMENT (\$ 542.00)

Complete if Known	
Application Number	
Filing Date	October 25, 2000
First Named Inventor	Peterson, Thomas A.
Examiner Name	
Group Art Unit	
Attorney Docket No.	P04716US2 (ISURF 2330)

METHOD OF PAYMENT

1. The Commissioner is hereby authorized to charge indicated fees and credit any overpayments to:

Deposit Account Number **26-0084**
 Deposit Account Name **Zarley, McKee, Thomte, Voorhees & Sease**

Charge Any Additional Fee Required
Under 37 CFR 1.16 and 1.17

Applicant claims small entity status.
See 37 CFR 1.27

2. Payment Enclosed:

Check Credit card Money Order Other

FEE CALCULATION

1. BASIC FILING FEE

Large Entity Small Entity

Fee Code (\$)	Fee Code (\$)	Fee Description	Fee Paid	
101	710	201	355 Utility filing fee	355
106	320	206	160 Design filing fee	
107	490	207	245 Plant filing fee	
108	710	208	355 Reissue filing fee	
114	150	214	75 Provisional filing fee	

SUBTOTAL (1) (\$ 355.00)

2. EXTRA CLAIM FEES

Total Claims	Extra Claims	Fee from below	Fee Paid
23	-20** = 3	X 9	= 27
Independent Claims	7	- 3** = 4	X 40 = 160
Multiple Dependent			

Large Entity Small Entity

Fee Code (\$)	Fee Code (\$)	Fee Description
103	18	203 9 Claims in excess of 20
102	80	202 40 Independent claims in excess of 3
104	270	204 135 Multiple dependent claim, if not paid
109	80	209 40 ** Reissue independent claims over original patent
110	18	210 9 ** Reissue claims in excess of 20 and over original patent

SUBTOTAL (2) (\$ 187.00)

**or number previously paid, if greater; For Reissues, see above

3. ADDITIONAL FEES


Large Entity Fee Code (\$)	Small Entity Fee Code (\$)	Fee Description	Fee Paid
105	130	205 65 Surcharge - late filing fee or oath	
127	50	227 25 Surcharge - late provisional filing fee or cover sheet	
139	130	139 130 Non-English specification	
147	2,520	147 2,520 For filing a request for ex parte reexamination	
112	920*	112 920* Requesting publication of SIR prior to Examiner action	
113	1,840*	113 1,840* Requesting publication of SIR after Examiner action	
115	110	215 55 Extension for reply within first month	
116	390	216 195 Extension for reply within second month	
117	890	217 445 Extension for reply within third month	
118	1,390	218 695 Extension for reply within fourth month	
128	1,890	228 945 Extension for reply within fifth month	
119	310	219 155 Notice of Appeal	
120	310	220 155 Filing a brief in support of an appeal	
121	270	221 135 Request for oral hearing	
138	1,510	138 1,510 Petition to institute a public use proceeding	
140	110	240 55 Petition to revive - unavoidable	
141	1,240	241 620 Petition to revive - unintentional	
142	1,240	242 620 Utility issue fee (or reissue)	
143	440	243 220 Design issue fee	
144	600	244 300 Plant issue fee	
122	130	122 130 Petitions to the Commissioner	
123	50	123 50 Petitions related to provisional applications	
126	240	126 240 Submission of Information Disclosure Stmt	
581	40	581 40 Recording each patent assignment per property (times number of properties)	
146	710	246 355 Filing a submission after final rejection (37 CFR § 1.129(a))	
149	710	249 355 For each additional invention to be examined (37 CFR § 1.129(b))	
179	710	279 355 Request for Continued Examination (RCE)	
169	900	169 900 Request for expedited examination of a design application	

Other fee (specify) _____

* Reduced by Basic Filing Fee Paid

SUBTOTAL (3) (\$)

SUBMITTED BY

Name (Print/Type)	Heidi S. Nebel	Registration No. (Attorney/Agent)	37,719	Telephone	515-288-3667
Signature					
Date	10/25/00				

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

JC914 U.S. PRO
09/696600
10/25/00



PATENT
P04716US2
ISURF 2330

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICANT: **PETERSON, Thomas A., et al.** ART UNIT:
SERIAL NO: **NEW APPLICATION** EXAMINER:
FILED:
TITLE: **METHODS AND MATERIALS TO INDUCE RECOMBINATION
IN PLANTS**

**STATEMENT TO SUPPORT FILINGS AND SUBMISSIONS
IN ACCORDANCE WITH 37 CFR §§1.821 - 1.825**

Commissioner of Patents and Trademarks
Washington, D.C. 20231

Dear Sir:


I, Heidi S. Nebel, hereby declare and state the
following:

1. I am the attorney of record in the above-identified case.
2. The content of the paper and computer readable copies of the sequence listing, submitted in accordance with 37 CFR §1.821(c) and (e) respectfully, are the same.

=====

CERTIFICATE OF MAILING (37 C.F.R. § 1.8(a))

I hereby certify that this **DECLARATION TO SUPPORT FILINGS AND SUBMISSIONS IN ACCORDANCE WITH 37 CFR §§1.821 - 1.825** has been mailed by U.S. Postal Service's "Post Office to Addressee" Express Mail service in an envelope addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231, Box Patent Application, prior to 5:00 p.m. on the 25 day of October, 2000.




Patricia E. Wilson
Express Mail label EL515383044US


I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under §1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application of any patent issued thereon.

Dated

10/25/00


Heidi S. Nebel

Respectfully submitted,


Heidi S. Nebel

Reg. No. 37,719

ZARLEY, McKEE, THOMTE, VOORHEES
& SEASE
ATTORNEYS OF RECORD

801 Grand Avenue, Suite 3200
Des Moines, Iowa 50309-2721
515-288-3667 -pw-

Applicant or Patentee: Thomas A. Peterson and Yongli Xiao
Serial No. or Patent No:
Filed or Issued: October 25, 2000
For: METHODS AND MATERIALS TO INDUCE RECOMBINATION IN PLANTS

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
37 CFR 1.9(f) AND 1.27(d) - NONPROFIT ORGANIZATION

I hereby declare that I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF ORGANIZATION Iowa State University Research Foundation, Inc.
ADDRESS OF ORGANIZATION 310 Lab of Mechanics
Ames, Iowa 50011-2131

TYPE OF ORGANIZATION

[] UNIVERSITY OR OTHER INSTITUTION OF HIGHER EDUCATION
[] TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 USC 501(a) and 501(c)(3))
[X] NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE

UNITED STATES OF AMERICA

(NAME OF STATE Iowa

(CITATION OF STATUTE Chapter 504A, Code of Iowa)

[] WOULD QUALIFY AS TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE
(26 USC 501(a) and 501(c)(3)) IF LOCATED IN THE UNITED STATES OF AMERICA
[] WOULD QUALIFY AS NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE
OF THE UNITED STATES OF AMERICA IF LOCATED IN THE UNITED STATES OF AMERICA
(NAME OF STATE _____)
(CITATION OF STATUTE _____)

I hereby declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9(e) for purposes of paying reduced fees under Section 41(a) and (b) of Title 35, United States Code with regard to the invention entitled METHODS AND MATERIALS TO INDUCE RECOMBINATION IN PLANTS by inventor(s) Thomas A. Peterson and Yongli Xiao as described in

the specification filed herewith.
 application serial no. _____, filed _____.
 patent no. _____, issued _____.

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above identified invention.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(d) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

* NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27).

FULL NAME _____
ADDRESS _____
 INDIVIDUAL SMALL BUSINESS CONCERN NONPROFIT ORGANIZATION

FULL NAME _____
ADDRESS _____
 INDIVIDUAL SMALL BUSINESS CONCERN NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of payment, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b)).
I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING Dr. Kenneth Kirkland
TITLE IN ORGANIZATION Executive Director, Iowa State University Research Foundation, Inc.
ADDRESS OF PERSON SIGNING 310 Lab of Mechanics, Ames, Iowa 50011-2131

SIGNATURE K. Kirkland DATE October 25, 2000

INVENTORS: Thomas Peterson
Yongli Xiao

TITLE: Methods and Materials to Induce Recombination
in Plants

Address correspondence to:

Heidi S. Nebel
Reg. No. 37,719
ZARLEY, McKEE, THOMTE,
VOORHEES & SEASE
801 Grand Suite 3200
Des Moines, Iowa 50309-2721
515-288-3667

TO ALL WHOM IT MAY CONCERN:

Be it known that we, Thomas Peterson and Yongli Xiao have invented certain new and useful improvements in **Methods and Materials to Induce Recombination in Plants** of which the following is a specification:

FIELD OF THE INVENTION

This application claims priority to Provisional Patent Application US 60/069,057 filed December 10, 1997.

BACKGROUND OF THE INVENTION

Transposons, or transposable elements, are mobile genetic elements characterized by their ability to move in and out of their host genome in a manner similar to bacteriophages. However, unlike bacteriophages, transposable elements are unable to move from host to host, since they lack the structural features necessary to package and deliver the nucleic acid.

Transposable elements have been shown to induce recombination in yeast (Boeke, ed. Berg & Howe, *Mobile DNA* 335 (Am. Soc. Microbio. 1989); drosophila (Davis et al., 84 PNAS 174 (1987) and Busseau et al., 218 *Mol. Gen. Genet.* 222

DRAFT 6/26/2018

(1989)) and plants (Martin et al., 119 *Genetics* 171 (1988), McClintock, 53 *Washington Year Book* 254 (1954), Doring et al., 184 *Mol. Gen. Genet.* 377 (1981), Taylor and Walbot, 4 *EMBO J.* (1985), Dooner et al., 211 *Mol. Gen. Genet.* 485 (1988); Lowe et al., 132 *Genetics* 813 (1992) and Athma and Peterson, 128 *Genetics* 163 (1991)). The recombinations resulted in deletions, duplications and other rearrangements. These studies were limited to observations of natural phenomenon in the organisms, and did not involve *in situ* recombination of non-self constructs.

Human manipulation of transposons has been limited, until the present invention, to insertion and excision of transposons, with or without accompanying nucleic acid. For example, US Patent 5,482,852 to Yoder et al. describes a technique for inserting an expression-cassette-containing transposon into plant cells, selecting cells which contain the transposon, and subsequently crossing the plant so as to remove the transposon sequences, but not the expression cassette. US Patent 5,013,658 to Dooner describes a method to use selectable markers in transposons as tags for genetic studies. US Patent 5,478,369 to Albertson describes a male-fertility mediating gene which was cloned through the use of the Ac (maize) transposon.

Human manipulation of recombination has been limited, until the present invention, to non-transposon related methods. US Patent 5,527,695 to Hodges et al. describes the use of a recombinase enzyme to induce recombination *in situ* of a construct with homologous direct repeats. US Patent 5,658,772 to Odell et al. describes a similar method, but the crossover site and the recombinase enzyme are more defined (lox and Cre, respectively). In *Arabidopsis*, generation of double strand breaks by HO endonuclease increased the frequency of somatic intrachromosomal homologous recombination by about 10 fold. Chiurazzi et al., 8 *Plant Cell* 2057 (1996). Moreover, double strand breaks are known to initiate recombination in fungi. Szostak et al., 33 *Cell* 25 (1983).

Citation of the above documents is not intended as an admission that any of the foregoing is pertinent prior art. All statements as to the date or representation as to the contents of these documents is based on subjective characterization of information available to the applicant at the time of writing, and does not constitute an admission as to the accuracy of the dates or contents of these documents.

SUMMARY OF THE INVENTION

It is therefore an object of the present invention to provide a molecular system useful to induce recombination in plants.

It is a further object to provide a molecular system useful to increase the frequency of recombination events in plants over the naturally-occurring frequency.

It is yet another object to provide a molecular system for introducing part or parts of a construct into a plant cell and subsequently causing the part or parts to recombine.

It is yet another object to provide a molecular system for inducing recombination in plants at particular times, in inserted sequences.

In all of the above embodiments, it is an object to provide a molecular system which is useful to direct timing of transcription of constructs and timing of expression of construct products.

In another embodiment, it is an object to provide a method to affect the timing of complementary nucleic acid production in a plant cell.

It is another object to provide a molecular system which is useful to affect naturally-occurring sequences.

It is also an object of the invention to provide an *in situ* molecular system for constructing fusion protein-encoding sequences as well as expression of the fusion protein encoded.

It is a further object to provide a molecular system for making directed change(s) in a construct after it has been integrated into the genome of a plant.

In all of these embodiments, it is an object to provide compositions useful to induce homologous recombination in plants, including constructs, plasmids, viruses, cells and plants.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is schematic of a GU-Ds-US construct.

Figure 2, upper panel, is a map of a GU-Ds-US construct, showing primer construct location and, lower panel, is a agarose gel showing PCR amplification products for certain pairs of primers.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to the unexpected finding that overlapping foreign gene sequences containing a maize Ds element can be induced to undergo homologous recombination upon introduction of the maize Ac transposase. The studies in this regard were in the *Arabidopsis* plant, using the overlapping, but incomplete, fragments of the GUS gene (a bacterial gene) to flank a Ds element. The Ac transposase was shown to induce recombination of the foreign GUS gene fragments to produce a functional GUS gene. The frequency of homologous recombination was increased by over 1000 fold over that which is normally seen in nature. This high recombination rate suggests that the action of the Ac transposase may not be explained solely by induction of a double strand break, but probably reflects a heretofore unknown effect of the Ac transposase.

Previous to the present invention, transposons were thought to be useful as passive vectors for inserting and deleting foreign constructs into and out of cells. In the present invention, transposons are viewed in a completely

different light; transposons, or their parts, can now be considered active tools for recombining foreign constructs *in situ*. Moreover, the Ac transposase can now be used to regulate the timing of functional transcripts and construct products without the need for inducible promoters or enhancers at the site of insertion.

Transposon-induced recombination can be useful to enhance the frequency of certain homology-dependent gene manipulations in plants. The *Arabidopsis* and maize genomes are currently being saturated with Ac/Ds insertions, and it is possible to induce recombination between endogenous sequence duplications that carry Ac/Ds insertions. Such recombination events can generate interstitial deletions that could simplify the analysis of multicopy gene clusters such as are frequently associated with disease resistance genes. Transposon-induced recombination can also be used to delete undesirable sequences from integrated transgenes, or to delete multiple transgene copies that are commonly generated in many transformation protocols.

Therefore, one broad aspect of the present invention is drawn to methods to induce homologous recombination in a plant, comprising introducing a recombination construct to a plant and making available to the plant a transposase, so as to induce recombination. Preferably, the recombination construct and transposase are from maize.

For example, the above aspect can be implemented in the following manners:

A. The recombination construct may be a simple Ds element which is transformed into a plant and inserted between known direct repeats in the genome.

B. The recombination construct may be a Ds element flanked by direct repeats. In one such embodiment, this construct would enable disruption of a gene and subsequent regeneration of function via homologous recombination.

C. The recombination construct may be a Ds element flanked by overlapping sequences having homologous regions.

In one such embodiment, the sequences, when recombined, may encode ie. a gene, coding region or fusion protein.

D. The recombination construct may be a Ds element-gene construct which is flanked by direct repeats. In one such embodiment, the direct repeats can be nonsense repeats, and in another, the direct repeats can encode a gene product when recombined.

All of the above recombination constructs may optionally contain a selection marker or other sequences.

E. The transposase can exist as a natural product of the plant, and the construct can be introduced to the plant which contains the transposase.

F. The transposase can be cotransformed, or present on the recombination construct, with or without an inducible promoter.

G. The transposase can be provided by way of a cross between a recombination construct-containing plant and a transposase-containing plant.

In all of the above aspects, the recombination construct and/or the transposase may be transiently or stably transformed into the plant. In this respect, the recombination construct or transposase genes may or may not be integrated into the genome of the plant. In all of the above aspects, the orientation of the Ds element and/or construct is not essential; either can be oriented in any manner. However, in all of the above aspects, it is preferable that the Ds element is between the direct repeats (for example, as in Fig. 1), although it is possible to induce recombination when the Ds element is within the direct repeats.

By "transposase" it is meant any plant transposase, ie. an enzyme that can induce homologous recombination in the presence of a Ds or Ac element.

By "Ac element" it is meant that which is typically understood in the art as the definition: any form of maize Ac, including any isolate or any allelic variant or homologue thereof.

By "plant" it is meant one or more plant seed, plant embryo, plant part or whole plant. The plant may be a monocot or dicot, man-made or naturally-occurring.

By "recombination construct" it is meant a full or partial (recombination-inducing) isolated Ds element either in conjunction with, or without, other isolated nucleic acid sequences, such as, for example, a regulatory sequence (promoter, enhancer, terminator, etc.), an isolated full gene sequence, an isolated coding region, or an isolated partial natural or artificial sequence. Therefore, an Ac element with altered transposase function is within this definition.

By "overlapping sequences having homologous regions" or "overlapping sequences having homologous sequences" it is meant that the entire sequence which encodes the construct product (or regulatory region) desired must be represented by the two fragments, and that the some of the same internal sequences must exist on each of the two fragments so as to allow recombination of the fragments to produce a complete construct. The gene product need not be a naturally-occurring gene product, nor need encode an independently-functional gene product. It could, for example, encode a fusion protein or sub-unit of a functional complex. Additionally, it could encode an complementary transcript useful for inhibiting translation of mRNA. Lastly, the sequences could be nonsense sequences used simply for their ability to recombine.

By "proteins" it is meant any compounds which comprise amino acids, including peptides, polypeptides, fusion proteins, etc.

By "Ds element" it is meant not only the definition known in the art, ie. a transposon or part of a transposon without the sequence for a functional transposase, but also the instance where the element possesses a transposase coupled with a regulatory region which is post-insertion manipulatable. For instance, it is within the present definition of "Ds element" the instance where the element is

an isolated natural transposon, with an inducible promotor inserted at the appropriate site upstream from the transposase, so as to allow human manipulation of the timing of transcription and expression of the transposase. However, the full transposon is not necessary. In other words, so long as the Ds element retains the ability to induce recombination, the presence or absence of sequences present in the naturally-occurring Ds element or transposon is not critical. "Ds element" includes the classic "dissociation"-type of element (ie, "state 1" and "state 2"), as well as the full gamut of more recently characterized Ds elements, ie. those with inverted repeats.

By "making available the transposase" it is meant that the transposase becomes available at levels to cause an increase in homologous recombination over that seen in the plant under normal conditions. In other words, the naturally-occurring Ac transposase, in the case of maize, could be caused to over-express, or a construct could be introduced which operably links an inducible promoter or enhancer to the Ac sequence. In all plants, the transposase can be introduced via co-transformation (on the same or separate vectors) with the recombination construct, or subsequent to the transformation of the recombination construct. In either case (co-transformation or subsequent transformation) the transposase can be linked operably to an inducible promoter or enhancer which would cause the gene product to be available in the cell at a pre-determined time. In some instances, it may be desirable to cause the transposase to be available as soon as the recombination construct has been introduced, and in such case, it may be desirable to introduce the vector containing the transposase prior to introduction of the recombination construct, so that the enzyme can be active as soon as possible subsequent to the introduction of the recombination construct. In other instances, it may be desirable to provide the transposase in such a manner that the enzyme would be induced upon a certain environmental or internal occurrence.

Another means for subsequently making available the Ac transposase is as is described in the Examples: crossing a plant transformed with the recombination construct with a plant expressing the transposase. The transposase in this instance can be naturally-occurring or provided via genetic engineering.

By "introducing" it is meant any means of transferring nucleic acid into the cell. For example, any of the following known means can be used: infective, vector-containing bacterial strains (such as *Agrobacterium rhizogenes* and *Agrobacterium tumefaciens*) according to ie. Zambryski, 43 *Ann. Rev. Pl. Physiol. Pl. Mol. Biol.* 465 (1992); pollen-tube transformation [Zhon-xun et al., 6 *Plant Molec. Bio.* 165 (1988)]; direct transformation of germinating seeds [Toepfer et al., 1 *Plant Cell* 133 (1989)]; polyethylene glycol or electroporation transformation [Christou et al., 84 *Proc. Nat. Acad. Sci.* 3662 (1987)]; and biolistic processes [Yang & Cristou, *Particle Bombardment Technology for Gene Transfer* (1994)].

The transformed cells may be induced to form transformed plants via organogenesis or embryogenesis, according to the procedures of Dixon *Plant Cell Culture: A Practical Approach* (IRL Press, Oxford 1987).

Any seed, embryo, plant or plant part is amenable to the present techniques. Of course, the agronomically-significant seeds, embryos, plants or plant parts are preferred. Soybean, maize, sugar cane, beet, tobacco, wheat, barley, poppy, rape, sunflower, alfalfa, sorghum, rose, carnation, gerbera, carrot, tomato, lettuce, chicory, pepper, melon and cabbage are among the preferred seeds, embryos, plants or plant parts. Particularly preferred are: soybean, tobacco and maize seeds, embryos, plants or plant parts. However, *Arabidopsis* seeds, embryos, plants or plant parts are also preferred, since it is an excellent system for study of plant genetics.

In the case of a recombination construct containing the Ds element and other isolated nucleic acid sequences, the

nucleic acid sequences can be any that are useful in plant genetics. Preferred are those genes or sequences which are agronomically significant. For example, genes encoding male sterility, foreign organism resistance (viruses or bacteria), including genes which produce bacterial endotoxins, such as *bacillus thuringiensis* endotoxin, genes involved in specific biosynthetic pathways (eg. in fruit ripening, oil or pigment biosynthesis, seed formation, or starch metabolism) or genes involved in environmental tolerance (eg. salt tolerance, drought tolerance, or tolerance to anaerobic conditions). The choice of gene or sequence induced to recombine in the present invention is not limited. Examples of genes and how to obtain them are available through reference articles, books and supply catalogs, such as *The Sourcebook* (1-800-551-5291). Sambrook et al., *Molecular Cloning. A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) and Weising et al., 22 *Ann Rev. Gen.* 421 (1988) also have pertinent information.

Plants which have undergone induced homologous recombination can be identified by Southern blot analysis, by identification of the particular gene product expressed (marker rescue) or deleted (marker loss) through recombination, or by the methods described in US Patent 5,482,852 to Yoder et al., (PCR, regain of function) which patent is herein incorporated by reference. These techniques are well-known in the art.

Where and when the sequences are inserted can also contribute to the overall desirability of the end product. For example, it may be desirable to select for seeds, embryos, or plant parts (all within the definition of "plants") which have incorporated the sequences in strategically-beneficial locations. In one instance, it may be desirable to select a plant which has incorporated a sequence so as to cause a disruption or other alteration of a coding or regulatory region of a genomic gene sequence. This is within skill of the art, and considered part of the present invention. Moreover, it may be desirable to

introduce the recombination construct and/or transposase gene to an undifferentiated cell or callus of cells, so that the entire plant contains the constructs and/or genes.

In the case of a recombination construct containing direct repeats, the direct repeats may be any length which allows homologous recombination to be induced upon introduction of the transposase. In general, the larger the length of the direct repeats the more likely it is that homologous recombination will be successful, although at certain long lengths (greater than 10 kilobases, perhaps) the relative increase over that seen with shorter lengths may be diminishing. In one embodiment of the present invention, the direct repeats are from 300 base pairs to 10 kilobase pairs in length, although it is possible to have fewer or more, depending on the particular application desired. Preferably, the direct repeats are 600 base pairs to 6 kilo base pairs in length.

In the case of a recombination construct containing direct repeats, the distance between the direct repeats (not including the length of the direct repeats themselves) may be any length which allows homologous recombination to be induced upon introduction of the transposase. In one embodiment of the present invention, the distance between the direct repeats is from approximately 25 base pairs to approximately 15 kilo base pairs. Preferably, the distance between the direct repeats is 6 kilobases to 12 kilobases, and includes an approximately 4.6 kilobase sequence of a Ds element.

Therefore, there are provided methods to induce transcription and/or translation of a gene in a plant comprising introducing to the plant a Ds element containing overlapping sequences having homologous regions, which sequences, when homologously recombined, result in a gene and subsequently making available to the plant the transposase, so as to induce homologous recombination and subsequent transcription and translation of said gene.

Preferred plants, genes and sequences are those described previously.

Also provided are methods to disrupt or otherwise alter a naturally-occurring sequence (including genes) in plants, comprising introducing to the plant a recombination construct having direct repeats useful for subsequent removal of the recombination construct via homologous recombination. Preferred plants, genes and sequences are those described previously. Transgenic knockout plants or plants with specific genomic changes can be obtained by these methods.

Moreover, methods to construct a fusion protein sequence in plants, comprising introducing to the plant a recombination construct having overlapping sequences having homologous regions, which sequences, when homologously recombined, result in a fusion protein sequence, and making available to the plant a transposase, so as to cause recombination and construction of a fusion protein sequence. Preferred plants, genes and sequence are those described previously.

Also provided is a method to induce complementary nucleic acid sequence production in plants, comprising introducing to the plant a recombination construct having overlapping sequences having homologous regions, which sequences, when homologously recombined, result in complementary nucleic acid sequences, and making available to the plant a transposase, so as to cause recombination and production of said complementary nucleic acid sequences. Preferred plants and genes are those described previously.

The present invention also provides materials useful in the present invention.

In particular, plants comprising a recombination construct are provided. Such plants are those which have been previously described, although maize and arabidopsis are preferred plants herein provided. The presently-identified plants can inherently possess the transposase activity, or have artificially modified transposase

activity. Plants wherein the transposase activity is provided via an inducible promoter is preferred, although plants which are crosses between a Ds-construct-transformed plant and a plant in which transposase is naturally occurring is also preferred.

Vector comprising the transposase and recombination construct are therefore also provided by the present invention. Vectors with either an early or late promoter in conjunction with a transposase and/or a recombination construct are particularly provided. Vectors may be from bacterial, viral or artificial origin. Vectors of the present invention optionally comprise the following promoters useful in early expression of plant sequences:

Ogs4B (Tsuchiya et al., 36 Plant Cell Physiology 487 (1994)

TA29 (Koltunow et al., 2 Plant Cell 1201 (1990)

A3 & A9 (Paul et al., 19 Plant Molecular Biology 611 (1992)

In order to then constitutively express the sequences described above, the vector optionally contains, for example, a 35S promoter.

Vectors may be obtained from various commercial sources, including Clontech Laboratories, Inc. (Palo Alto, CA), Stratagene (La Jolla, CA), Invitrogen (Carlsbad, CA), New England Biolabs (Beverly, MA) and Promega (Madison, WI).

Genes can be obtained through catalogues such as *The Sourcebook* (1-800-551-5291 or conventional methods as in Sambrook et al., *Molecular Cloning. A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989)

The Ac transposase sequence can be obtained by Genbank Accession Number: X01380, and either cloned according to Sambrook (above) or obtained as has been previously described.

A Ds element can be obtained by using the Ac element (above), or portion thereof as a probe to detect a Ds element or Ac element. If the element retrieved is an Ac element, internal sequences can be removed so as to

eliminate transposase function. Alternatively, because the terminal inverted repeats of Ds (and Ac) elements are conserved, and the subterminal sequences are somewhat conserved, primers can be designed and used for PCR amplification of a Ds element or Ac element. Again, if an Ac element is amplified, internal sequences can be removed so as to eliminate transposase function, so as to obtain a Ds element. Of course, an Ac element can also be modified via insertion of an -inducible promoter designed to drive the transposase. Oligonucleotide primers that can be used to PCR amplify Ac or Ds elements from maize are 5'-CAGGGATGAAAGTAG-3' (SEQ ID NO 1, from the 5' end of Ac terminal sequence) and 5'-TAGGGATGAAAACGG-3' (SEQ ID NO 2, from the 3' end of Ac terminal sequence). The methods for conducting these types of molecular manipulations are well-known in the art, and are described in detail in Sambrook et al., *Molecular Cloning. A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) and Ausubel et al., *Current Protocols in Molecular Biology* (Greene Publishing Associates, Inc., 1993). Moreover, Ds elements can be obtained from researchers willing to send samples, or from other publicly-available sources. For instance, the Arabidopsis Biological Resource Center, Ohio State University, 309 B&Z Building, 1735 Neil Avenue, Columbus, OH 43210, (Phone: 614-292-9371) will provide plasmid pWS31 (stock number CD3-125), containing DsE, and plasmid pWS32 (stock number CD3-126) containing DsG. Other Ds elements can be obtained from the American Type Culture Collection, Rockville, MD (800) 638-6597.

Direct repeats can be synthesized de novo, or retrieved from plasmid preparations carrying the desired sequences. For instance, if overlapping sequences having homologous regions are desired, a plasmid preparation can be divided into two batches, with one batch subjected to a restriction enzyme digestion which cleaves at one site, and the other batch subjected to a restriction enzyme digestion which cleaves at a different, overlap-causing site. The methods

for conducting these types of molecular manipulations are well-known in the art, and are described in detail in Sambrook *et al.*, *Molecular Cloning. A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) and Ausubel *et al.*, *Current Protocols in Molecular Biology* (Greene Publishing Associates, Inc., 1993).

Recombination constructs can be made using the starting materials above or with additional materials, using methods well-known in the art. In general, the sequences can be manipulated to have ligase-compatible ends, and incubated with ligase to generate full constructs. For example, restriction enzymes can be chosen on the basis of their ability to cut at an acceptable site in both sequence to be ligated, or a linker may be added to convert the sequence end(s) to ones that are compatible. The methods for conducting these types of molecular manipulations are well-known in the art, and are described in detail in Sambrook et al., *Molecular Cloning. A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) and Ausubel et al., *Current Protocols in Molecular Biology* (Greene Publishing Associates, Inc., 1993). The methods described herein according to Tinland et al., 91 Proc. Natl. Acad. Sci. USA 8000 (1994) can also be used.

Constructs with Ac transposase can be made according to the methods described in Calvi et al., 66 *Cell* 465 (1991); Long et al., 241 *Mol. Genet.* 627 (1993); and Honma et al., 90 *Proc. Natl. Acad. Sci.* 6242 (1993).

The present invention also includes compositions of matter which can be induced to undergo homologous recombination upon introduction of a maize transposase comprising a maize recombination construct having proximal direct repeat sequences. By "proximal" it is meant that the Ds element is flanked, either with or without intervening sequences, by the direct repeats. That is, so long as the Ds element is internal to at least two direct repeats, such that at least one direct repeat is at either end of the Ds element, the direct repeat sequences are "proximal". Other

PCT/US2006/008660

sequences can exist between the direct repeats as well. The direct repeats can be artificial or naturally-occurring, isolated sequences. The recombination construct can be made according to the methods detailed previously.

Also provided are compositions of matter which can be induced to undergo homologous recombination upon introduction of a maize transposase comprising a maize recombination construct having proximal direct repeat sequences as part of a vector. The vector can be any nucleic acid sequence or other carrier (ie. liposomes) useful for transferring the recombination construct during molecular manipulations. These compositions can be made according to the methods detailed previously.

Also provided are compositions of matter which can be induced to undergo homologous recombination upon introduction of a maize transposase comprising a maize recombination construct having proximal direct repeat sequences in a plant. These compositions can be made using any plants, especially the plants disclosed herein as useful in the present invention. Compositions as described, which further comprise a transposase gene under control of an inducible promoter, or a gene internal to said direct repeat sequences, or wherein the direct repeat sequences are in the form of overlapping sequences having homologous regions are preferred. These compositions can be made according to the methods detailed previously.

The following are examples are not intended to limit the scope of the present invention as described and claimed. They are simply for the purpose of illustration.

Examples

Example 1 - Preparation of nucleic acids

A Ds element (courtesy of V. Sundaresan) was inserted between two partially overlapping non-functional segments of the (-glucuronidase gene5 (GUS; courtesy of H. Puchta) to

generate a binary plant transformation vector termed GU-Ds-US (Figure 1). The sequences were prepared according to Tinland et al., 91 *Proc. Natl. Acad. Sci. USA* 8000 (1994) and the plasmid containing the sequences was called "plasmid GU.US." The homologous direct repeat sequences are 618 bp in length, and the distance between them is 6.3 kb including the 4.6 kb Ds element.

In particular, plasmid pWS31 (described in Sundaresan et al., 9 *Genes Dev.* 1797 (1995)) with a Ds element was cut with Sal I and religated to remove the GUS gene in pWS31 (according to Sambrook et al., *Molecular Cloning. A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989)), and the new plasmid was named pWS31Y. SacI was used to digest pWS31Y, and the 4.8 kb Ds fragment was purified via an 0.8% agarose gel using Promega Wizard DNA Clean-up System according to Ausubel et al., *Current Protocols in Molecular Biology* (Greene Publishing Associates, Inc., 1993).

Plasmid GU.US was digested with PstI restriction enzyme and Promega Wizard DNA Clean-Up System was used to purify the DNA, as above. A PstI-SacI adaptor was designed, and subsequently synthesized by ISU DNA Synthesis Facility, using standard synthesis procedures.

The PstI cut GU.US, the 4.8kb Ds fragment and the PstI-SacI adaptor were ligated together, and subsequently transformed into *E.coli*. A plasmid GU-Ds-US was obtained. Sequence from the cloning junction confirmed proper sequence configuration for further studies.

Example 2 - Transformation of plants

The electroporation method described in Ausubel et al., *Current Protocols in Molecular Biology* (Greene Publishing Associates, Inc., 1993) was used to transform GU-Ds-US into Agrobacterium strain ASE. The vacuum infiltration method was used to transform Agrobacterium into Arabidopsis, Columbia ectotype. Seeds were selected on the plate with

30ug/ml kanamycin. After one week, transformants were transferred from selection plates to soil. Plants were grown in growth chamber under conditions of 16 hours light/8 hours dark at 20-22 centigrade. These procedures were according to Ellis et al., 316 C. R. Acad. Sci. Paris 1194 (1993); Bent et al. 265 Science 1856 (1994) and Koncz et al., 204 Mol. Gen. Genet. 383 (1986) .

Example 3 - Growth of plants/identification of recombination

The T1 seeds were harvested from the plants transformed according to Example 2. From each independent transformant, 17 seeds were planted. Southern hybridization was used to check the insertion number of the GU-Ds-US construct to get the single insertion transformant. Then, the T2 seeds were harvested from single insertion lines and the seeds were put on selection plates to obtain homozygous lines of the transgene. The plants with the single insertion and the plants homozygous for the transgene were used as the starter lines.

The starter lines and lines with stable Ac transposase (obtained from Arabidopsis Stock Center (CS8045, CS8537, CS8538) were planted. The starter lines were used as the pollen donor and the stable Ac lines as the pollen receptor in crosses. At the same time, the starter line pollen was crossed to wild-type(WT) Arabidopsis (No-O and Landsberg) as control.

F1 heterozygous seeds (GU-Ds-US/-, Ac- and GU-Ds-US/-) were planted. When the plants were at full rosette stage, the whole plant was used for X-Glu staining according to Jefferson, 5 Plant. Mol. Biol. Rep. 387 (1987) and Swoboda et al., 13 EMBO J. 484 (1994).

Simple excision of Ds was predicted to not restore an intact GUS gene, whereas homologous recombination between the partially-duplicated GUS segments would give a functional GUS gene detectable by staining. Comparisons between sibling plants with and without Ac transposase

indicates the degree to which homologous recombination is stimulated by Ac transposase over the level of spontaneous recombination. The results show that Ac induces recombination of the GU-Ds US substrate at a surprisingly high frequency in Arabidopsis.

As shown in Table 1, plants in the control group (genotype GU-Ds-US/+; no Ac) had zero or few blue sectors (average of 0.6 blue sectors per plant), whereas plants in the experimental group commonly had hundreds of blue sectors (average of 700 blue sectors per plant). These results were obtained from three independently transformed GU-Ds-US lines (DsI5, DsI6, DsII7), in crosses with three different lines expressing Ac transposase (CS8045, CS8537, CS8538). Based on the average numbers of blue sectors, the calculated spontaneous recombination frequency of the GU-Ds-US transgene in the absence of Ac transposase is approximately 1.99×10^{-7} to 5.40×10^{-8} events/genome3. This frequency is similar to that reported by others for spontaneous recombination of a similar GU.US transgene^{3,7,8}. In the presence of the sAc transposase source, the average recombination frequency is increased >1,000 fold.

The data from the above experiment is as follows:

Table 1: Frequency of recombination in GU-Ds-US-transformed Arabidopsis plants, with and without available Ac transposase. Recombination is shown by blue spots (in CS 8045 x Ds I 5).

Type of Plant	Number of Plants	Total Number of Blue Spots
Control: Gu-Ds-US	9	3
Experimental: GU-Ds-US + Transposase	10	3,310

Table 2: Frequency of recombination in GU-Ds-US-transformed Arabidopsis plant parts in the presence of Ac transposase. Recombination is shown by blue spots per part (in CS 8045 x Ds I 5).

Plant	Cotyled on (parts/ plant)	Leaf (parts/ plant)	Stem	Root	Total
1	15 (2)	348 (9)	9	43	415
2	33 (2)	18 (8)	4	16	71
3	50 (2)	419 (10)	5	62	536
4	17 (2)	298 (10)	0	59	374
5	22 (2)	397 (10)	0	57	476
6	21 (2)	90 (10)	9	63	174
7	20 (2)	259 (9)	6	64	349
8	35 (2)	222 (10)	5	23	285
9	5 (1)	169 (7)	2	25	201
10	61 (2)	324 (7)	8	36	429

Example 4 - PCR and Southern Confirmation of recombination construct recombination

Genomic DNA samples were amplified by PCR for 35 cycles as follows: denaturation at 94°C for 30 sec; annealing at 57°C for 30 sec; elongation for 1 min at 72°C for 35 cycles. PCR products were cloned into pT7Blue T-vector (Novagen) and sequenced at the Iowa State University Nucleic Acid Facility, Ames , IA. Probe preparations and Southern blot hybridizations were performed as described.

The structure of the GU-Ds-US transgene in the presence or absence of the Ac transposase source (sAc) using PCR amplification of genomic DNA was determined. A band predicted to arise from the restored GUS gene was obtained using DNA from plants containing the GU-Ds-US construct together with sAc, but not from plants lacking sAc. The PCR band from the presumptive recombination product was sequenced and found to contain a precisely restored GUS gene. Similarly, a PCR product predicted to arise from Ds excision from the GU-Ds-US construct was found only in plants containing sAc, but not in plants lacking sAc. The sequence of the presumptive Ds excision product showed that it contained a typical Ds excision footprint.

Among the progeny of the variegated GUS+ plants, was detected several plants with uniform GUS+ expression. To test for germlinal transmission of a recombined GUS transgene, the genomic DNA from plants with uniform GUS expression by Southern hybridization was analyzed. Hybridization with a GUS-specific probe detected two bands (7.3 kb and 4.7 kb) in HindIII-digested DNA from a plant of genotype GU-Ds-US/-, sAc/- (Figure 4a, lane 2), whereas the GUS-specific probe detected only one band of 5.0 kb in DNA from two plants with uniform GUS expression. These results are expected from the generation of a single GUS coding sequence by recombination between the homologous regions of the GU-Ds-US construct. To verify that the recombinant GUS+ plants were derived from the original GU-Ds-US transformants, additional Southern hybridizations were performed using enzymes that cleave in the DNA flanking the transgene insertion. Genomic DNA from plants of genotype GU-Ds-US, sAc and the uniform GUS+ plants were digested with EcoRV and hybridized with a probe specific for the 3' end of the GU-Ds-US construct (Figure 1). The probe hybridizes to the same 4.2 kb band in the GU-Ds-US progenitor plants and the uniform GUS+ plants. This result indicates that the two GUS+ plants could not have arisen by seed or pollen contamination, but did in fact originate by recombination of the GU-Ds-US transgene locus.

Although the present invention has been fully described herein, it is to be noted that various changes and modifications are apparent to those skilled in the art. Such changes and modifications are to be understood as included within the scope of the present invention as defined by the appended claims.

WHAT IS CLAIMED IS:

1. A method to induce homologous recombination in a plant, comprising introducing a recombination construct to the plant, and making available to the plant a transposase, so as to induce homologous recombination.
2. A method of claim 1, wherein the recombination construct comprises a maize Ds element and the transposase is of maize origin.
3. A method of claim 2, wherein the recombination construct further comprises direct repeats proximal to the Ds element, and an agronomically significant gene internal to the direct repeats, wherein the agronomically significant gene is selected from the group consisting of: genes useful for disease resistance; genes useful for male sterility; genes useful for environmental condition tolerance; and genes useful for commercially-enhancing a biosynthetic pathway.
4. A method of claim 2, wherein the plant in which recombination is induced is selected from the group consisting of: soybean; maize; sugar cane; beet; tobacco; wheat; barley; poppy; rape; sunflower; alfalfa; sorghum; rose; carnation; gerbera; carrot; tomato; lettuce; chicory; pepper; melon; *Arabidopsis*; and cabbage.
5. A method to construct a functional gene in plants, comprising introducing to the plant a maize recombination construct having overlapping sequences having homologous regions, which sequences, when homologously recombined, result in a functional gene, and making available to the plant a maize transposase, so as to induce recombination and construction of a functional gene.

6. A method of claim 5, wherein the functional gene is selected from the group consisting of: genes useful for disease resistance; genes useful for male sterility; genes useful for environmental condition tolerance; and genes useful for commercially-enhancing a biosynthetic pathway.

7. A method of claim 5, wherein the plant in which recombination is induced is selected from the group consisting of: soybean; maize; sugar cane; beet; tobacco; wheat; barley; poppy; rape; sunflower; alfalfa; sorghum; rose; carnation; gerbera; carrot; tomato; lettuce; chicory; pepper; melon; *Arabidopsis*; and cabbage.

8. A method to induce transcription and/or translation of a gene in a plant comprising introducing to the plant a maize Ds element containing overlapping sequences having homologous regions, which sequences, when homologously recombined, result in a gene; and subsequently making available to the plant the transposase, so as to induce homologous recombination and subsequent transcription and translation of said gene.

9. A method of claim 8, wherein the gene is selected from the group consisting of those genes useful for: genes useful for disease resistance; genes useful for male sterility; genes useful for environmental condition tolerance; and genes useful for commercially-enhancing a biosynthetic pathway.

10. A method of claim 9, wherein the plant in which recombination is induced is selected from the group consisting of: soybean; maize; sugar cane; beet; tobacco; wheat; barley; poppy; rape; sunflower; alfalfa;

sorghum; rose; carnation; gerbera; carrot; tomato; lettuce; chicory; pepper; melon; *Arabidopsis*; and cabbage.

11. A method to alter a naturally-occurring sequence in plants, comprising introducing to the plant a maize recombination construct having direct repeats useful for subsequent removal of the recombination construct via homologous recombination.

12. A method of claim 11, wherein the plant in which the naturally-occurring sequence is disrupted is selected from the group consisting of: soybean; maize; sugar cane; beet; tobacco; wheat; barley; poppy; rape; sunflower; alfalfa; sorghum; rose; carnation; gerbera; carrot; tomato; lettuce; chicory; pepper; melon; *Arabidopsis*; and cabbage.

13. A method to construct a fusion protein sequence in plants, comprising introducing to the plant a maize recombination construct having overlapping sequences having homologous regions, which sequences, when homologously recombined, result in a fusion protein sequence, and making available to the plant a maize transposase, so as to cause recombination and construction of a fusion protein sequence.

14. A method of claim 2, wherein the fusion protein sequence is selected from the group consisting of: genes useful for disease resistance; genes useful for male sterility; genes useful for environmental condition tolerance; and genes useful for commercially-enhancing a biosynthetic pathway.

15. A method of claim 13, wherein the plant in which a fusion protein is constructed is selected from the group consisting of: soybean; maize; sugar cane; beet;

DESEMBOLGADO

tobacco; wheat; barley; poppy; rape; sunflower; alfalfa; sorghum; rose; carnation; gerbera; carrot; tomato; lettuce; chicory; pepper; melon; Arabidopsis; and cabbage.

16. A method to induce complementary nucleic acid sequence production in plants, comprising introducing to the plant a maize recombination construct having overlapping sequences having homologous regions, which sequences, when homologously recombined, result in complementary nucleic acid sequences, and making available to the plant a maize transposase, so as to cause recombination and production of said complementary nucleic acid sequences.

17. A method of claim 16, wherein the plant in which recombination and production of complementary nucleic acid sequences is induced is selected from the group consisting of: soybean; maize; sugar cane; beet; tobacco; wheat; barley; poppy; rape; sunflower; alfalfa; sorghum; rose; carnation; gerbera; carrot; tomato; lettuce; chicory; pepper; melon; Arabidopsis; and cabbage.

18. A composition of matter which can be induced to undergo homologous recombination upon introduction of a maize transposase comprising a maize recombination construct having proximal direct repeat sequences.

19. A composition of matter which can be induced to undergo homologous recombination upon introduction of a maize transposase comprising a maize recombination construct of claim 18 as part of a vector.

20. A composition of matter which can be induced to undergo homologous recombination upon introduction of a

maize transposase comprising a maize recombination construct of claim 18 in a plant.

21. A composition of matter of claim 20, which further comprises a transposase gene under control of an inducible promoter.

22. A composition of matter of claim 20, which further comprises a gene internal to said direct repeat sequences.

23. A composition of matter of claim 20, wherein said direct repeat sequences are in the form of overlapping sequences having homologous regions.

ABSTRACT OF THE DISCLOSURE

The present invention is drawn to methods and materials to induce homologous recombination in a plant, comprising introducing a recombination construct to a plant and making available to the plant a transposase, so as to induce recombination. Preferably, the recombination construct and transposase are from maize.

DRAFT - PLEASE DO NOT CITE






FIGURE 1



PCR primers pairs:

1. Negative control
- 2, 7. H2+Ds(-)
- 3, 8. GUS1+H3
- 4, 9. H1+H2
- 5, 10. 35S+(U)S
- 6, 11. Ac primers,
Ac6+AcX6

FIGURE 2

SEQUENCE LISTING

<110> Peterson, Thomas
Xiao, Yongli

<120> Methods and Materials to Induce Recombination in Plants

<130> P-1033

<140> filed herewith
<141> 1998-12-09

<150> 60/069057

<151> 1997-12-10

<160> 2

<170> PatentIn Ver. 2.0

<210> 1
<211> 15
<212> DNA
<213> Zea mays

<400> 1
cagggatgaa agtag

15

<210> 2
<211> 15
<212> DNA
<213> Zea mays

<400> 2
tagggatgaa aacgg

15

Sequence Listing

DECLARATION FOR PATENT APPLICATION AND APPOINTMENT OF ATTORNEY

As a below-named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name; I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention (Design, if applicable), entitled : Methods and Materials to Induce Recombination in Plants

the specification of which (check one):

is attached hereto.

was filed on _____ as Application Serial No. _____ and was amended on _____ (if applicable).

was filed on _____ as International Application (PCT) No. _____ and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with *Title 37, Code of Federal Regulations, § 1.56(a)*. I hereby claim foreign priority benefits under *Title 35, United States Code § 119* of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which the priority is claimed.

PRIOR FOREIGN APPLICATION(S)

NUMBER	COUNTRY	DAY/MONTH/YEAR FILED	PRIORITY CLAIMED
			Yes No
			Yes No

I hereby claim the benefit under *Title 35, United States Code, § 120* of any United States application(s) or PCT international application(s) designating The United States of America listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of *Title 35, United States Code, § 112*, I acknowledge the duty to disclose material information as defined in *Title 37, Code of Federal Regulations, § 1.56(a)* which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

APPLICATION NUMBER	FILING DATE	STATUS (Patented, Pending or Abandoned)
60/069,057	December 10, 1997	Pending


I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine, or imprisonment, or both, under *Section 101 of Title 18 of the United States Code*, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: I (We) hereby appoint as my (our) attorneys, with full powers of substitution and revocation, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith: Kristine H. Johnson, Registration Number 36,835, Jennifer Bales 38,070, and Jean Macheleit 33,956..

Send correspondence to: KRISTINE H. JOHNSON
THE OPERA GALLERIA, SUITE 219
123 NORTH COLLEGE AVENUE
FORT COLLINS, CO 80524

TELEPHONE CALLS TO: KRISTINE H. JOHNSON
(970) 472-9650

See following pages for additional joint inventors.

First Name of First or Sole Inventor Thomas A. Peterson	Citizenship USA		
Residence Address - Street 27947 560th Avenue	Post Office Address Street 27947 560th Avenue		
City Ames			
State or Country Iowa	Zip 50010	State or Country Iowa	Zip 50010
DATE 9 December 1998	SIGNATURE 		

DECLARATION FOR PATENT APPLICATION AND APPOINTMENT OF ATTORNEY

Page 2

Full Name of Additional Joint Inventor Yongli Xiong	Citizenship P.R. China		
Residence Address - Street 1256 Hawthorn Court	Post Office Address Street 1256 Hawthorn Court		
City Ames	City		
State or Country Iowa	Zip 50010	State or Country Iowa	Zip 50010
Date 12/09/98	Signature 		

Full Name of Additional Joint Inventor	Citizenship		
Residence Address - Street	Post Office Address Street		
City	City		
State or Country	Zip	State or Country	Zip
Date	Signature		

Full Name of Additional Joint Inventor	Citizenship		
Residence Address - Street	Post Office Address Street		
City	City		
State or Country	Zip	State or Country	Zip
Date	Signature		

Full Name of Additional Joint Inventor	Citizenship		
Residence Address - Street	Post Office Address Street		
City	City		
State or Country	Zip	State or Country	Zip
Date	Signature		

SEE FOLLOWING PAGES FOR ADDITIONAL JOINT INVENTORS.